

Revealed cores: characterizations and structure

Stefano Vannucci

Dipartimento di Economia Politica, Università di Siena
Piazza San Francesco 7, 53100 Siena (Italy)

September 3, 2013

Abstract

Characterizations of the choice functions that select the cores or the externally stable cores induced by an underlying revealed dominance digraph are provided. Relying on such characterizations, the basic order-theoretic structure of the corresponding sets of revealed cores is also analyzed. In particular, it is shown that the poset of all revealed cores ordered by set inclusion is a median meet semilattice: therefore, any profile of revealed cores may be aggregated by means of the simple majority rule.

JEL classification: C70, C71, D01, D03

Keywords: Core, choice functions, dominance digraphs, revealed preference

1 Introduction

The *core* of a game is the set of its *undominated* outcomes, with respect to a suitably defined *dominance* irreflexive relation, or loopless digraph. Now, consider the ongoing operation of a multi-agent system, e.g. an organization or indeed any decision-making unit whose outputs are aptly modeled as the outcomes of a game. Let us then assume that the set of available options does in fact change at a faster pace than the behavioural attitudes of the relevant players *and* the latter interact as predicted by the core of that game. It follows that the corresponding choice behaviour of the given interaction system as recorded by its choice function should be constrained in some way by its game-theoretic structure and thus somehow *reveal* that fact. But then, what are the characteristic ‘fingerprints’ of such a choice function, namely the *testable* behavioural predictions of the core as a solution concept? Or more simply, *which choice functions defined over arbitrary subsets of an ‘universal’ outcome set may be regarded as revealed cores?* Let us call that issue, for ease of reference, the (full domain) *core revelation problem*.

Apparently, such a problem has never been addressed in its *full generality* in the extant literature. To be sure, parts of the massive body of literature on ‘revealed preference’ provide partial answers addressing the case of *nonempty cores*, i.e. of *acyclic* revealed dominance digraphs (see e.g. Wilson (1970), Sen (1971), Plott (1974), Suzumura (1983)). Moreover, there is also some work covering the case of possibly empty sets of undominated outcomes for an *arbitrary* - i.e. possibly *not* irreflexive- binary relation R , hence putting aside the original game-theoretic interpretation of R as a dominance relation (see e.g. Aizerman and Aleskerov (1995), and Danilov and Koshevoy (2009)). But of course the dominance relation of a game in its usual meaning has to be *irreflexive* (no outcome dominates itself), and the core of a game may well be *empty*, because its *revealed dominance digraph may have cycles*. Here, we are interested precisely in the *general* version of the core revelation problem for the full domain, namely in *a characterization of all revealed cores as solutions for a certain ‘universal’ outcome set and all of its subsets, including (locally) empty-valued cores*.

A further generalization of the core revelation problem to the case of choice functions with an arbitrary domain (along the lines of Bossert, Suzumura (2010)) is left as a topic for another paper.

The present paper is aimed at filling this gap in the literature by addressing the general core revelation problem with full domain as formulated above,

under *several* variants of the notion of core. A study of the basic order-theoretic properties of the corresponding classes of revealed core-solutions as canonically ordered by set-inclusion is also provided: in particular, it is shown that the class of *all* revealed cores (as opposed to, say, the class of nonempty-valued revealed cores) is a *meet sub-semilattice* of the lattice of all choice functions, and in fact a *median meet semilattice*. A remarkable consequence of that fact is that *any profile of revealed cores is amenable to aggregation by the simple majority rule*.

Thus, it turns out that each revealed core embodies a considerable part of standard maximizing choice, while the global structure of (full domain) revealed cores retains the order-theoretic properties of the space of all (full domain) choice functions that is most significant from the point of view of simple majority aggregation.

The paper is organized as follows: section 2 includes a presentation of the model and the main characterization results; section 3 provides some basic results concerning the order-theoretic properties of the classes of revealed core-solutions previously characterized; section 4 consists of a few concluding remarks.

2 Choice functions and revealed cores

Let X be a set denoting the ‘universal’ outcome set, with cardinality $\#X \geq 3$, and $\mathcal{P}(X)$ its power set. It is also assumed for the sake of convenience that X is *finite* (but it should be remarked that the bulk of the ensuing analysis is easily lifted with suitable minor adaptations to the case of an infinite outcome set). A *choice function on X* (with full domain) is a *deflationary* operator on $\mathcal{P}(X)$ i.e. a function $c : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ such that $c(A) \subseteq A$ for any $A \subseteq X$ (empty choice sets are allowed). A choice function c is *proper* if $c(A) \neq \emptyset$ whenever $\emptyset \neq A \subseteq X$. We denote C_X the set of all choice functions on X , and C_X° the subset of all *proper* choice functions on X . The *proper subdomain* of $c \in C_X$ -written D_c - is the set of all subsets of X with a nonempty-valued choice set i.e. $D_c = \{A \subseteq X : c(A) \neq \emptyset\}$. For any binary relation $\mathcal{B} \subseteq X \times X$, and any $Y \subseteq X$, \mathcal{B}^a and \mathcal{B}^s denote the asymmetric and symmetric components of \mathcal{B} , respectively, while $\mathcal{B}_Y = \mathcal{B} \cap (Y \times Y)$ and $\overline{\mathcal{B}} = (X \times X) \setminus \mathcal{B}$. Recall that $\mathcal{B} \subseteq X \times X$ is *reflexive* iff $x\mathcal{B}x$ for all $x \in X$, *irreflexive* iff *not* $x\mathcal{B}x$ for all $x \in X$, *total* iff $x\mathcal{B}y$ or $y\mathcal{B}x$ for any $x, y \in X$, *asymmetric* iff $x\mathcal{B}y$ entails *not* $y\mathcal{B}x$ for any $x, y \in X$, *transitive* iff $x\mathcal{B}y$

and $y\mathcal{B}z$ entail $x\mathcal{B}z$ for any $x, y, z \in X$, *quasi-transitive* if \mathcal{B}^a is transitive, *negatively transitive* if $\overline{\mathcal{B}}$ is transitive. The transitive closure $T(\mathcal{B})$ is the smallest transitive $R \supseteq \mathcal{B}$. Moreover, \mathcal{B} is *strictly acyclic* iff its transitive closure is irreflexive, and a *strict partial order* iff it is both asymmetric and transitive.

Let $\Delta \subseteq X \times X$ be an *irreflexive* binary relation on X , denoting a suitably defined *dominance* relation: (X, Δ) is the corresponding *dominance digraph* (in graph-theoretic parlance, (X, Δ) is in particular a simple, loopless digraph i.e. a directed graph with at most one arc between any ordered pair of vertices, and with no arc from any vertex to itself). In particular, (X, Δ) is an *asymmetric dominance digraph* if $\Delta = \Delta^a$.

For any $Y \subseteq X$, $\Delta_Y = \Delta \cap (Y \times Y)$ denotes the dominance relation induced by Δ on Y (of course $\Delta_X = \Delta$), and (Y, Δ_Y) is the induced *dominance subdigraph* on Y . Broadly speaking, the *core* of (Y, Δ_Y) is the set of Δ_Y -undominated outcomes in Y , namely

$$\mathbb{C}(Y, \Delta_Y) = \{y \in Y : \text{not } z\Delta_Y y \text{ for all } z \in Y\}.$$

The *a-core* of (Y, Δ_Y) is the set of Δ_Y^a -undominated outcomes in Y , namely $\mathbb{C}^a(Y, \Delta_Y) = \mathbb{C}(Y, \Delta_Y^a) = \{y \in Y : \text{not } z(\Delta_Y)^a y \text{ for all } z \in Y\}$.

The *core* (*a-core*) of (Y, Δ_Y) is *externally stable* iff for any $z \in Y \setminus \mathbb{C}(Y, \Delta_Y)$ there exists $y \in \mathbb{C}(Y, \Delta_Y)$ such that $y\Delta_Y z$ (for any $z \in Y \setminus \mathbb{C}^a(Y, \Delta_Y)$ there exists $y \in \mathbb{C}^a(Y, \Delta_Y)$ such that $y(\Delta_Y)^a z$, respectively).

A dominance digraph (X, Δ) is also said to be *core-perfect* or *strictly acyclic* (*acyclic*, respectively) if $\mathbb{C}(Y, \Delta_Y) \neq \emptyset$ ($\mathbb{C}^a(Y, \Delta_Y) \neq \emptyset$, respectively) for any $Y \subseteq X$.

Remark 1 *It should be emphasized here that any dominance digraph may arise in a natural way from an underlying game in coalitional form and from a related game in strategic form. Indeed, the dominance digraph (X, Δ_g^{a*}) defined by the following rule can be attached in a natural way to any coalitional game $g = (N, X, E, (\succ_i)_{i \in N})$:*

for any $x, y \in X$, $x, y \in X$, $x\Delta_g^{a*} y$ iff there exist $A \subseteq X$ and $S \subseteq N$ such that

$x \in A \in E(S)$ and $z \succ_i y$ for all $i \in S$ and $z \in A$ (see Vannucci (2009) for further details).

Two binary relations $R(c)$, R_c induced by a choice function $c \in C_X$ on X and defined as follows will play a pivotal role in the ensuing analysis: for

any $x, y \in X$, $xR_c y$ if and only if $x \in c(\{x, y\})$, while $xR(c)y$ if and only if there exists $Y \subseteq X$ such that $x \in c(Y)$ and $y \in Y$.

A choice function $c \in C_X$ is a *revealed core-solution* if there exists an *irreflexive* relation $\Delta \subseteq X \times X$ such that $c(Y) = \mathbb{C}(Y, \Delta_Y)$ for any $Y \subseteq X$. Similarly, $c \in C_X$ is a *revealed a-core-solution* (*ES core-solution*, *ES a-core-solution*, respectively) if there exists an *irreflexive* relation $\Delta \subseteq X \times X$ such that $c(Y) = \mathbb{C}^a(Y, \Delta_Y)$ ($c(Y) = \mathbb{C}(Y, \Delta_Y)$ with $\mathbb{C}(Y, \Delta_Y)$ externally stable, $c(Y) = \mathbb{C}_u(Y, \Delta_Y)$, $c(Y) = \mathbb{C}_u^a(Y, \Delta_Y)$, respectively) for any $Y \subseteq X$. Then, we also say that c is *core-rationalizable* (*a-core-rationalizable*, *ES-core-rationalizable*, *ES-a-core-rationalizable* respectively) by the dominance digraph (X, Δ) . Clearly, *ES (a-)core-solutions* are refinements of *(a-)core solutions*. *Revealed cores* will also be used as a generic label to denote all the foregoing choice functions.

Example 2 Notice that digraph (X, \emptyset) is also a dominance digraph, and $\mathbb{C}(A, \emptyset_A) = \mathbb{C}^a(A, \emptyset_A) = A$ for any $A \subseteq X$ (hence it is also -trivially-externally stable). Therefore, the identity operator $c^{id} : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ is a revealed core-solution (*a-core-solution*, *ES core-solution*).

Example 3 By way of contrast, take $\emptyset \subseteq G \subset X$ and consider the dichotomic choice function $c_-^G : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ as defined by the ‘strict’ satisficing rule $c_-^G(A) = A \cap G$ for any $A \subseteq X$. It is easily checked that c_-^G is not a revealed core: to see this, take any $x \in X \setminus G$. Then, $c_-^G(\{x\}) = \emptyset$ while for any dominance digraph (X, Δ) and any $x \in X$, it cannot be the case that $x\Delta x$ hence $\mathbb{C}(\{x\}, \Delta_{\{x\}}) = \mathbb{C}^a(\{x\}, \Delta_{\{x\}}) = \{x\}$.

Example 4 Next, take again $\emptyset \subseteq G \subset X$ and consider the nonempty valued dichotomic choice function $c_+^G : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ as defined by the ‘lax’ satisficing rule $c_+^G(A) = A \cap G$ for any $A \subseteq X$ if $A \cap G \neq \emptyset$, and $c_+^G(A) = A$ otherwise. Now, posit $\Delta = G \times (X \setminus G)$ i.e. $x\Delta y$ iff $x \in G$ and $y \in X \setminus G$. It is easily checked that for any $Y \subseteq X$, $c_+^G(Y) = \mathbb{C}(Y, \Delta_Y) = \mathbb{C}^a(Y, \Delta_Y)$ (which is also externally stable).

The main objective of this article is precisely to provide a *characterization* of all revealed cores in C_X , and study their basic order-theoretic structure.

To begin with, let us consider two requirements concerning local existence of nonempty choice sets.

No-dummy property (ND): $c(\{x\}) = \{x\}$ for any $x \in X$.

2- Properness (2-PR): $c(A) \neq \emptyset$ for any $A \subseteq X$ such that $\#A = 2$.

It is easily checked that ND is satisfied by all revealed cores, while 2-PR is only violated by core solutions when the underlying dominance digraph is *not* asymmetric. A stronger property that obviously entails both ND and 2-PR is

Properness (PR): $c(A) \neq \emptyset$ for any nonempty $A \subseteq X$.

The following properties of a choice function $c \in C_X$ play a prominent role, under various labels, in the extant literature:

Chernoff Contraction-consistency (C): for any $A, B \subseteq X$ such that $A \subseteq B$, $c(B) \cap A \subseteq c(A)$.

Concordance (CO): for any $A, B \subseteq X$, $c(A) \cap c(B) \subseteq c(A \cup B)$.

Superset consistency (SS): for any $A, B \subseteq X$, if $A \subseteq B$ and $\emptyset \neq c(B) \subseteq c(A)$ then $c(A) \subseteq c(B)$.

Property C is a contraction-consistency condition for choice sets in that it requires that any outcome chosen out of a certain set should also be chosen out of any subset of the former: essentially, it says that *any good reason to choose a certain option out of a given menu should retain its strength in every submenu of the former containing that option*.

Conversely, property CO (also variously denoted as γ or Generalized Condorcet-consistency) is an expansion-consistency condition for choice sets, requiring that an outcome chosen out of a certain set and of a second one should also be chosen out of the larger set given by the union of those two sets: it says that *any good reason to choose a certain option out of two given menus should retain its strength in the larger menu obtained by merging those two menus*.

Property SS is also an expansion-consistency requirement for choice sets: it rules out the possibility that the choice set of a certain menu be nonempty *and* strictly included in the choice sets of a smaller menu.

We are now ready to prove the main results of this paper. Let us start from the following simple

Claim 5 Let $R \subseteq X \times X$ be any (binary) relation on X , and define $\Delta^R \subseteq X \times X$ by the following rule: for any $x, y \in X$, $x\Delta^R y$ iff $\text{not } yRx$. Then,

- (i) $\Delta^{\Delta^R} = R$;
- (ii) for any $Y \subseteq X$, $\max R_Y = \{x \in Y : \text{not } y\Delta^R x \text{ for all } y \in Y\}$, and $\max \Delta^R_Y = \{x \in Y : \text{not } yRx \text{ for all } y \in Y\}$;
- (iii) R is reflexive iff Δ^R is irreflexive, and irreflexive iff Δ^R is reflexive;
- (iv) R is total iff Δ^R is asymmetric, and asymmetric iff Δ^R is total;
- (v) R is quasi-transitive iff Δ^R is quasi-transitive.

Proof. (i) For any $x, y \in X$, by definition $x\Delta^{\Delta^R} y$ iff $\text{not } y\Delta^R x$ iff $\text{not } (\text{not } xRy)$ iff xRy .

(ii) Let $x \in Y$, and xRy for all $y \in Y$: then, by definition, $\text{not } y\Delta^R x$ for all $y \in Y$, and conversely if $\text{not } y\Delta^R x$ for all $y \in Y$ then $\text{not } (\text{not } xRy)$ i.e. xRy for all $y \in Y$. Similarly, $x \in Y$ and $\text{not } yRx$ for all $y \in Y$: then by definition $x\Delta^R y$ for all $y \in Y$, and conversely.

(iii) Indeed, by definition for any $x \in X$, $\text{not } x\Delta^R x$ iff $\text{not}(\text{not } xRx)$ i.e. xRx . Similarly, $\text{not } xRx$ iff $x\Delta^R x$.

(iv) Suppose Δ^R is asymmetric: then, for any $x, y \in X$, it may be the case that $\text{not } y\Delta^R x$ or $\text{not } x\Delta^R y$ (or both). Now, if $\text{not } y\Delta^R x$ then xRy and if $\text{not } x\Delta^R y$ then yRx , therefore R is total. Conversely, suppose R is total. If xRy then $\text{not}(\text{not } xRy)$ hence $\text{not}(y\Delta^R x)$ and similarly yRx entails $\text{not}(x\Delta^R y)$, thus in any case Δ^R is asymmetric. Similarly, R is asymmetric iff for any $x, y \in X$ it cannot be the case that xRy and yRx , i.e. by definition iff it is not the case that $\text{not } y\Delta^R x$ and $\text{not } x\Delta^R y$, namely Δ^R is total.

(v) Suppose that R is quasi-transitive, and that both $x(\Delta^R)^a y$ and $y(\Delta^R)^a z$. Then, by definition ($\text{not } yRx$ and xRy), and ($\text{not } zRy$ and yRz) i.e. $xR^a y$ and $yR^a z$, hence $xR^a z$. Therefore, xRz and $\text{not } zRx$ i.e. $\text{not } z\Delta^R x$ and $x\Delta^R z$, namely $x(\Delta^R)^a z$. Conversely, suppose that Δ^R is quasi-transitive, and that both $xR^a y$ and $yR^a z$. Then, by definition (xRy and $\text{not } yRx$), and (yRz and $\text{not } zRy$) i.e. by definition ($\text{not } y\Delta^R x$ and $x\Delta^R y$) and ($\text{not } z\Delta^R y$ and $y\Delta^R z$), i.e. $x(\Delta^R)^a y$ and $y(\Delta^R)^a z$, hence $x(\Delta^R)^a z$. Therefore, $x\Delta^R z$ and $\text{not } z\Delta^R x$ i.e. $\text{not } zRx$ and xRz , namely $xR^a z$. ■

Remark 6 The content of the previous Claim is certainly not unknown, but I have been unable to find a reference in print to it except for the statement of point (iv) in Monjardet (2007), while Plott (1974), Theorem 8, only includes a specialized version of the same point.

The following Theorem extends and/or supplements some previous characterization results for revealed cores due to Wilson (1970) and Sen (1971).

Theorem 7 *Let $c \in C_X$. Then, the following statements are equivalent:*

- (i) *c satisfies ND, C and CO;*
- (ii) *there exists a dominance digraph (X, Δ) such that $c(Y) = \mathbb{C}(Y, \Delta_Y)$ for any $Y \subseteq X$;*
- (iii) *there exists a reflexive relation $R \subseteq X \times X$ such that $c(Y) = \max R_Y$ for any $Y \subseteq X$.*
- (iv) *$R(c) = R_c$, $R(c)$ is reflexive and $c(Y) = \max R(c)_Y$ for any $Y \subseteq X$.*

Proof. (i) \implies (iv): Let $c \in C_X$. Now, for each $Y \subseteq X$ and $x \in c(Y)$, $xR(c)y$ for any $y \in Y$, by definition of $R(c)$. Hence $c(Y) \subseteq \max R(c)_Y$. Now, let $c \in C_X$ also satisfy ND, C and CO, and $x \in \max R(c)_Y$. Then, by definition, $x \in Y$ and for any $y \in Y$ there exists Y_y such that $y \in Y_y$ and $x \in c(Y_y)$. It follows that $x \in Y \subseteq \cup_{y \in Y} Y_y$ and, by CO, $x \in c(\cup_{y \in Y} Y_y)$ whence $x \in c(Y)$ by C. Therefore, $c(Y) = \max R(c)_Y$ (clearly it might be the case that $\max R(c)_Y = \emptyset$). Notice however that, by ND, $x \in c(\{x\})$ i.e. $xR(c)x$ for any $x \in X$. Thus, $R(c)$ is *reflexive*, as required. Moreover, if $xR(c)y$ then by C it must also be the case that $x \in c(\{x, y\})$ whence $xR_c y$ and thus $R(c) = R_c$ (since $R_c \subseteq R(c)$ by definition).

(ii) \iff (iii) (see Wilson (1970), Theorem 3): Let $c \in C_X$. Thus, by Claim 5 (ii), if there exists $R \subseteq X \times X$ such that $c(Y) = \max R_Y$ for any $Y \subseteq X$, then $c(Y) = \{x \in Y : \text{not } y\Delta^R x \text{ for all } y \in X\}$, for any $Y \subseteq X$. Moreover, if R is reflexive then by Claim 5(iii) Δ^R is irreflexive hence $c(Y) = \mathbb{C}(Y, \Delta_Y^R)$. Conversely if there exists an irreflexive $\Delta \subseteq X \times X$ such that $c(Y) = \mathbb{C}(Y, \Delta_Y)$ for any $Y \subseteq X$ then by Claim 5 (ii)-(iii) $c(Y) = \max \Delta_Y^R$ for any $Y \subseteq X$, and Δ^R is reflexive.

(iii) \implies (iv): See Wilson (1970), Theorem 3. Moreover, observe that $R_c \subseteq R(c)$ by definition, and $xR(c)y$ implies $x \in \max R(c)_{\{x, y\}} = c(\{x, y\})$ i.e. $xR_c y$ hence $R_c = R(c)$ (of course, this is an extension to arbitrary choice functions of the proof of the same result for *proper* choice functions due to Sen (1971)).

(iv) \implies (iii): Trivial.

(iii) \implies (i): Suppose that there exists a *reflexive* relation $R \subseteq X \times X$ such that $c(Y) = \max R_Y$ for any $Y \subseteq X$. Clearly, by reflexivity of R , $c(\{x\}) = \max R_{\{x\}} = \{x\}$, hence c satisfies ND. Moreover, for any $Y \subseteq Z \subseteq X$ and any $x \in c(Z) = \max R_Z$, it must also be the case that $x \in \max R_Y = c(Y)$

hence C is also satisfied by c . Finally, for any $Y, Z \subseteq X$ and $x \in X$, if $x \in c(Y) = \max R_Y$ and $c(Z) = \max R_Z$ then clearly $x \in \max R_{Y \cup Z}$ whence $x \in c(Y \cup Z)$ and CO is satisfied as well. ■

Remark 8 Notice that the equivalence between statements (ii) and (iii) of Theorem 7 above should in fact be credited to Wilson(1970) because it is strictly related (indeed, essentially equivalent) to Theorem 3 of Wilson (1970), though the latter concerns nonempty core-solutions over an arbitrary domain $D \subseteq \mathcal{P}(X) \setminus \{\emptyset\}$ hence, strictly speaking, is a statement about a class of partial proper choice functions. On the other hand, Aizerman and Aleskerov (1995) have a similar result (see their Theorem 2.5) namely a characterization by the conjunction of C and CO of the choice functions selecting the outcomes ‘permitted’ by all outcomes -or ‘not prohibited’ by any outcome- according to an arbitrary ‘permission’ or ‘prohibition’ binary relation.

Remark 9 The foregoing characterization result is tight. To check that, consider the following examples.

(1) Let $c^I \in C_X$ be defined as follows: for any $A \subseteq X$, $c^I(A) = \max L_A$ if $A \neq \{x^*\}$, and $c^I(\{x^*\}) = \emptyset$ where L is a linear order on X and x^* is its bottom element. Clearly, c^I violates ND , but satisfies C and CO ;

(2) Let $X = \{x, y, z\}$, and $c^{II} \in C_X$ be defined as follows: $c^{II}(\{h\}) = \{h\}$ for any $h \in X$, $c^{II}(\{x, y\}) = \{x\}$, $c^{II}(\{y, z\}) = \{y\}$, $c^{II}(\{x, z\}) = \{z\}$, and $c^{II}(X) = X$. It is immediately checked that c^{II} satisfies ND and CO , but violates C since e.g. $y \in c^{II}(X) \cap \{x, y\}$ but $y \notin c^{II}(\{x, y\})$;

(3) Let $c^{III} \in C_X$ be defined as follows: for any $A \subseteq X$, $c^{III}(A) = \max L_A$ if $\#A \leq 2$ and $c^{III}(A) = \emptyset$ otherwise, where L is a linear order on X . It is easily seen that c^{III} satisfies ND and C , but violates CO .

Next, we have a similar characterization result for revealed a-cores which is also an extension to the general case of possibly non-proper choice functions of previous results as discussed below (see Remark 13).

Theorem 10 Let $c \in C_X$. Then, the following statements are equivalent:

- (i) c satisfies ND , $2-PR$, C and CO ;
- (ii) there exists a dominance digraph (X, Δ) such that $c(Y) = \mathbb{C}^a(Y, \Delta_Y)$ for any $Y \subseteq X$;
- (iii) there exists a total relation $R \subseteq X \times X$ such that $c(Y) = \max R_Y$ for any $Y \subseteq X$;
- (iv) $R(c) = R_c$, $R(c)$ is total and $c(Y) = \max R(c)_Y$ for any $Y \subseteq X$.

Proof. (i) \implies (iii): Let $c \in C_X$ satisfy ND, 2-PR, C, and CO. Then, by ND, C and CO (and in view of Theorem 7 above) there exists a reflexive relation R on X such that $c(Y) = \max R_Y = \{y \in Y : yRz \text{ for all } z \in Y\}$ for each $Y \subseteq X$. Thus, by 2-PR, R is total.

(ii) \iff (iii): Suppose that there exists a total relation $R \subseteq X \times X$ such that $c(Y) = \max R_Y$ for any $Y \subseteq X$. Then, as recorded by Claim 5(ii) $c(Y) = \{x \in Y : \text{not } y\Delta^R x \text{ for all } y \in X\}$ for any $Y \subseteq X$. By Claim 5 (iv) Δ^R is asymmetric since R is total, hence in particular $c(Y) = \mathbb{C}(Y, \Delta_Y^R) = \mathbb{C}^a(Y, \Delta_Y^R)$ for any $Y \subseteq X$. Conversely, suppose that there exists a dominance digraph (X, Δ) such that $c(Y) = \mathbb{C}^a(Y, \Delta_Y) = \{x \in Y : \text{not } y\Delta^a x \text{ for all } y \in X\}$, for any $Y \subseteq X$. Then, as recorded by Claim 5 (ii) $c(Y) = \max(\Delta_Y^R)$: by Claim 5 (iv), $(\Delta^a)^R$ is total since Δ^a is asymmetric.

(ii) \implies (i): Suppose that there exists a dominance digraph (X, Δ) such that $c(Y) = \mathbb{C}^a(Y, \Delta_Y)$ for any $Y \subseteq X$. For any $x \in X$, $\text{not } x\Delta_{\{x\}}x$ i.e. $\text{not } x\Delta x$ by irreflexivity of Δ whence by definition $c(\{x\}) = \mathbb{C}^a(\{x\}, \Delta_{\{x\}}) = \mathbb{C}(\{x\}, \Delta_{\{x\}}) = \{x\}$ and ND is therefore satisfied by c . Furthermore, for any $x, y \in X$, $\Delta_{\{x,y\}} \subseteq \{(x, y), (y, x)\}$ hence $\Delta_{\{x,y\}}^a \in \{\emptyset, \{(x, y)\}, \{(y, x)\}\}$. If $\Delta_{\{x,y\}}^a = \emptyset$ then $\mathbb{C}^a(\{x, y\}, \Delta_{\{x,y\}}) = \{x, y\}$, otherwise $\mathbb{C}^a(\{x, y\}, \Delta_{\{x,y\}}) = \{x\}$ or $\mathbb{C}^a(\{x, y\}, \Delta_{\{x,y\}}) = \{y\}$, respectively, hence in any case

$$c(\{x, y\}) = \mathbb{C}^a(\{x, y\}, \Delta_{\{x,y\}}) \neq \emptyset \text{ thus } c \text{ satisfies 2-PR.}$$

Also, for any $Y, Z \subseteq X$ such that $Y \subseteq Z$, and any $x \in c(Z) \cap Y = \mathbb{C}^a(Z, \Delta_Z) \cap Y$, it must be the case that $\text{not } z\Delta_Z^a x$ for all $z \in Z$ hence in particular $\text{not } z\Delta_Y^a x$ for all $z \in Y$, i.e. $x \in \mathbb{C}^a(Y, \Delta_Y) = c(Y)$ and c also satisfies C.

Moreover, let $Y, Z \subseteq X$ and $x \in c(Y) \cap c(Z) = \mathbb{C}^a(Y, \Delta_Y) \cap \mathbb{C}^a(Z, \Delta_Z)$. Then, by definition, $\text{not } y\Delta_Y^a x$ for all $y \in Y$ and $\text{not } z\Delta_Z^a x$ for all $z \in Z$ hence $\text{not } u\Delta_{Y \cup Z}^a x$ for all $u \in Y \cup Z$ i.e. $x \in \mathbb{C}^a(Y \cup Z, \Delta_{Y \cup Z}) = c(Y \cup Z)$ and CO is satisfied by c .

(iii) \iff (iv): See the proof of Theorem 7 above. ■

Remark 11 *The foregoing characterization result is also tight. To see this, consider the following examples.*

(1) Let $c^I \in C_X$ as defined above (see Remark 9). Clearly, c^I violates ND, but satisfies 2-PR, C and CO;

(2) Let $c^{II^*} \in C_X$ be defined as follows: $c^{II^*}(\{x\}) = \{x\}$ for any $x \in X$, and $c^{II^*}(A) = \emptyset$ for any $A \subseteq X$ such that $\#A \geq 2$. It is easily checked that c^{II^*} does indeed satisfy ND, C and CO, but clearly violates 2-PR;

- (3) Let $X = \{x, y, z\}$, and $c^{II} \in C_X$ as defined above (see Remark 9). It is immediately checked that c^{II} satisfies ND, 2-PR, and CO, but violates C;
- (4) Let $c^{III} \in C_X$ as defined above (see Remark 9). It is easily seen that c^{III} satisfies ND, 2-PR and C, but violates CO.

Corollary 12 (see also Sen (1971), Suzumura (1983)) Let $c \in C_X^\circ$. Then, the following statements are equivalent:

- (i) c satisfies C and CO;
- (ii) there exists a strictly acyclic dominance digraph (X, Δ) such that $c(Y) = \mathbb{C}(Y, \Delta_Y) = \mathbb{C}^a(Y, \Delta_Y)$ for any $Y \subseteq X$;
- (iii) there exists a total relation $R \subseteq X \times X$ such that $c(Y) = \max R_Y$ for any $Y \subseteq X$;
- (iv) there exists a relation $R \subseteq X \times X$ such that $c(Y) = \max R_Y$ for any $Y \subseteq X$.
- (v) $R(c) = R_c$, $R(c)$ is total, and $c(Y) = \max R(c)_Y$ for any $Y \subseteq X$.

Proof. (i) \implies (ii): Since $c \in C_X^\circ$, c is proper hence in particular it also satisfies ND and 2-PR. Therefore, by Theorem 10 (ii) above, there exists a dominance digraph (X, Δ) such that $c(Y) = \mathbb{C}^a(Y, \Delta_Y)$ for any $Y \subseteq X$. Moreover, since by hypothesis c is *proper*, $\mathbb{C}^a(Y, \Delta_Y) \neq \emptyset$ for any $Y \subseteq X$ hence (X, Δ) must be *acyclic*. In particular, $\mathbb{C}^a(\{x, y\}, \Delta_{\{x, y\}}) \neq \emptyset$ for any $x, y \in X$, therefore Δ is asymmetric as well. Thus, (X, Δ) is indeed *strictly acyclic* and $\mathbb{C}(Y, \Delta_Y) = \mathbb{C}^a(Y, \Delta_Y) \neq \emptyset$ for any $Y \subseteq X$.

(ii) \implies (i): See the proof of Theorem 7 above;

(i) \iff (iii): Obvious, by Theorem 10 above, since, again, $c \in C_X^\circ$ entails that c satisfies ND and 2-PR.

(iii) \iff (iv): Suppose there exists $R \subseteq X \times X$ such that $c(Y) = \max R_Y$ for any $Y \subseteq X$. Since $c \in C_X^\circ$, $c(Y) \neq \emptyset$ for any $Y \subseteq X$. Hence, in particular, for any $x, y \in X$, $c(\{x, y\}) \neq \emptyset$. It follows that R is total. The reverse implication is trivial.

(iii) \iff (v): See the proof of Theorem 6 above, and of course Sen (1971).

■

Remark 13 Actually, it is well-known that a proper c satisfies both C and CO if and only if there exists a binary relation R on X such that $c(Y) = \max R_Y = \{y \in Y : yRz \text{ for all } z \in Y\}$ for each $Y \subseteq X$ and, moreover, $R = R(c) = R_c$ as defined above -indeed, $R(c) = R_c$ for any choice function that

satisfies C (see e.g. Sen (1971), Suzumura (1983)). Also notice that the equivalence between (ii) and (iii) is due to Plott (1974). Thus, Corollary 12 is -essentially- a restatement of the Sen-Plott-Suzumura characterization of revealed ‘rational’ (proper) choice functions or, equivalently, revealed non-empty core solutions.

Let us now turn to characterizations of *revealed externally stable core-solutions*. Since externally stable cores (of nonempty sets) are nonempty the corresponding choice functions are proper: thus, given the traditional focus on proper choice functions, this subclass of revealed cores is the most widely studied, and best known (thanks again to Wilson (1970) and Suzumura (1983)). Therefore, for the sake of convenience, we collect in the following Theorem a few notable characterizations of revealed externally stable cores (only some of them are already known and available in print, namely those recorded in Suzumura(1983) as explicitly mentioned in the proof below).

Theorem 14 *Let $c \in C_X$. Then, the following statements are equivalent:*

- (i) c satisfies PR, C , CO and SS;
- (ii) there exists a quasi-transitive relation $R \subseteq X \times X$ such that $c(Y) = \max R_Y \neq \emptyset$ for any nonempty $Y \subseteq X$;
- (iii) there exists a total and quasi-transitive relation $R \subseteq X \times X$ such that $c(Y) = \max R_Y \neq \emptyset$ for any nonempty $Y \subseteq X$;
- (iv) $R(c) = R_c$, $R(c)$ is total and quasi-transitive, and $c(Y) = \max R(c)_Y \neq \emptyset$ for any $Y \subseteq X$.
- (v) there exists a reflexive and negatively transitive relation $R \subseteq X \times X$ such that $c(Y) = \max R_Y \neq \emptyset$ for any nonempty $Y \subseteq X$;
- (vi) there exists a negatively transitive relation $R \subseteq X \times X$ such that $c(Y) = \max R_Y \neq \emptyset$ for any nonempty $Y \subseteq X$;
- (vii) there exists a dominance digraph (X, Δ) such that $c(Y) = \mathbb{C}(Y, \Delta_Y) = \mathbb{C}^a(Y, \Delta_Y)$ with $\mathbb{C}(Y, \Delta_Y)$ externally stable, for any $Y \subseteq X$;
- (viii) there exists a dominance digraph (X, Δ) such that Δ is transitive and $c(Y) = \mathbb{C}(Y, \Delta_Y) = \mathbb{C}^a(Y, \Delta_Y) \neq \emptyset$ for any nonempty $Y \subseteq X$;
- (ix) there exists a dominance digraph (X, Δ) such that Δ is a strict partial order and $c(Y) = \mathbb{C}(Y, \Delta_Y) = \mathbb{C}^a(Y, \Delta_Y) \neq \emptyset$ for any nonempty $Y \subseteq X$.

Proof. (i) \implies (ii) (Suzumura (1983)): By Theorem 2.6 of Suzumura (1983), if c satisfies PR, C, CO and SS then there exists a (*reflexive and*) quasi-transitive relation $R \subseteq X \times X$ such that $c(Y) = \max R_Y \neq \emptyset$ for any nonempty $Y \subseteq X$. But of course PR entails that $c(\{x, y\}) = \max R_{\{x, y\}} \neq \emptyset$ for any $x, y \in X$, hence R is *total* as well.

(ii) \implies (i) (Suzumura (1983)): See again Suzumura (1983), Theorems 2.5, 2.6 and 2.7.

(ii) \iff (iii): Let be $R \subseteq X \times X$ quasi-transitive and such that $c(Y) = \max R_Y \neq \emptyset$ for any nonempty $Y \subseteq X$. Of course, PR entails that in particular $c(\{x, y\}) = \max R_{\{x, y\}} \neq \emptyset$ for any $x, y \in X$, hence R is *total* as well. The reverse implication is trivial.

(iii) \iff (iv): See the proof of Theorem 7 above.

(iii) \iff (v): Let $R \subseteq X \times X$ be total and quasi-transitive, and $x, y, z \in X$ such that *not* xRy and *not* yRz . Hence, yRx and zRy since R is total. Therefore, by definition, yR^ax and zR^ay . By quasi-transitivity, it follows that zR^ax , whence in particular *not* xRz i.e. R is negatively transitive. Moreover, totality implies reflexivity of R . Conversely, let $R \subseteq X \times X$ be reflexive and negatively transitive. Suppose there exist $x, y \in X$ such that *not* xRy and *not* yRx : then, by negative transitivity, *not* xRx , a contradiction since R is reflexive. Thus, R is also total. Moreover, let xR^ay and yR^az . Then, in particular, *not* yRx and *not* zRy . It follows that, by negative transitivity, *not* zRx whence, by totality, xRz . Thus, xR^az i.e. R is quasi-transitive as well.

(v) \iff (vi): Let $R \subseteq X \times X$ be a *negatively transitive* relation such that $c(Y) = \max R_Y \neq \emptyset$ for any nonempty $Y \subseteq X$. Then in particular, $c(\{x\}) = \max R_{\{x\}} \neq \emptyset$ for any $x \in X$, hence R is reflexive as well. The reverse implication is trivial.

(iii) \implies (vii): Let be $R \subseteq X \times X$ total, quasi-transitive and such that $c(Y) = \max R_Y \neq \emptyset$ for any nonempty $Y \subseteq X$. Clearly, by construction, $c(Y) = \{x \in Y : xRy \text{ for all } y \in Y\}$ i.e.

$c(Y) = \{x \in Y : \text{not } y\Delta^R x \text{ for all } y \in Y\} = \mathbb{C}(Y, \Delta_Y^R)$ for any $Y \subseteq X$ (see Claim 5 (i) above). Moreover, by Claim 5 (iii), Δ^R is asymmetric since R is total, hence $\mathbb{C}(Y, \Delta_Y^R) = \mathbb{C}^a(Y, \Delta_Y^R)$. Now, take any $y_1 \in Y \setminus \mathbb{C}(Y, \Delta_Y^R)$. By definition, there exists $y_2 \in Y$ such that $y_2\Delta_Y^R y_1$. If $y_2 \in \mathbb{C}(Y, \Delta_Y^R)$ we are done. Suppose then that $y_2 \in Y \setminus \mathbb{C}(Y, \Delta_Y^R)$ as well: thus, there exists $y_3 \in Y$ such that $y_3\Delta_Y^R y_2$. It follows, by finiteness of Y and nonemptiness of $\mathbb{C}(Y, \Delta_Y^R)$, that there exists a finite k such that $y_i\Delta_Y^R y_{i-1}$ for any $i = 2, \dots, k$, and $y_k \in \mathbb{C}(Y, \Delta_Y^R)$. Since Δ^R is asymmetric, it also follows that $y_k\Delta_Y^R y_1$,

hence $\mathbb{C}(Y, \Delta_Y^R)$ is externally stable.

(vii) \implies (i): Suppose that there exists a dominance digraph (X, Δ) such that $c(Y) = \mathbb{C}(Y, \Delta_Y) = \mathbb{C}^a(Y, \Delta_Y)$ with $\mathbb{C}(Y, \Delta_Y)$ externally stable, for any $Y \subseteq X$. By definition of external stability, $c(Y) \neq \emptyset$ for any nonempty $Y \subseteq X$, hence c satisfies PR. Moreover, by Theorem 7 (ii) above (or, for that matter, by Theorem 8 (ii)), it also satisfies C and CO. Finally, consider $Y \subseteq Z \subseteq X$ such that $c(Z) \subseteq c(Y)$, and suppose there exists $y \in c(Y) \setminus c(Z)$ i.e. $y \in \mathbb{C}(Y, \Delta_Y) \setminus \mathbb{C}(Z, \Delta_Z)$. Then, by external stability of $\mathbb{C}(Z, \Delta_Z)$, there exists $z \in \mathbb{C}(Z, \Delta_Z) \subseteq \mathbb{C}(Y, \Delta_Y) \subseteq Y$ such that $z\Delta y$, a contradiction since $y \in \mathbb{C}(Y, \Delta_Y)$. Therefore, c satisfies SS as well.

(viii) \iff (iii): Suppose that there exists a dominance digraph (X, Δ) such that Δ is transitive (hence in particular quasi-transitive) and $c(Y) = \mathbb{C}(Y, \Delta_Y) = \mathbb{C}^a(Y, \Delta_Y) \neq \emptyset$ for any nonempty $Y \subseteq X$. Then, by Claim 5 (i)-(ii) above, $\emptyset \neq c(Y) = \mathbb{C}(Y, \Delta_Y) = \mathbb{C}(Y, \Delta_Y^{R^\Delta}) = \max R_Y^\Delta$ for any nonempty $Y \subseteq X$. Moreover, by Claim 5 (v), R^Δ is quasi-transitive. Also, notice that since by hypothesis Δ is both irreflexive and transitive, it must be *asymmetric* as well. Therefore, by Claim 5 (iv), R^Δ is total. Conversely, suppose that there exists a total and quasi-transitive relation $R \subseteq X \times X$ such that $c(Y) = \max R_Y \neq \emptyset$ for any nonempty $Y \subseteq X$. Then, by Claim 5 (ii) $c(Y) = \max R_Y = \mathbb{C}_u(Y, \Delta_Y^R) \neq \emptyset$ for any nonempty $Y \subseteq X$. Moreover, by Claim 5 (iii),(v), and in view of quasi-transitivity and totality of R , Δ^R is both quasi-transitive and asymmetric, hence *transitive* as well, and such that $\mathbb{C}_u(Y, \Delta_Y^R) = \mathbb{C}_u^a(Y, \Delta_Y^R)$ as required.

(viii) \iff (ix): Suppose that there exists a dominance digraph (X, Δ) such that Δ is *transitive* and $c(Y) = \mathbb{C}(Y, \Delta_Y) = \mathbb{C}^a(Y, \Delta_Y) \neq \emptyset$ for any nonempty $Y \subseteq X$. Again, irreflexivity and transitivity imply *asymmetry* of Δ , which is therefore a *strict partial order*. The reverse implication is trivial. ■

Remark 15 *Observe that the characterization result of revealed externally stable cores in terms of properties of choice functions included in Theorem 14 is also tight. To see this, consider the following examples.*

(1) Let $c^I \in C_X$ as defined above (see Remark 9). Clearly, c^I violates PR, but satisfies C, CO and SS;

(2) Let $X = \{x, y, z\}$, and $c^{II} \in C_X$ as defined above (see Remark 9). It is immediately checked that c^{II} satisfies PR, CO and SS, but violates C;

(3) Let $X = \{x, y, z\}$, and $c^{IV} \in C_X$ such that $c^{IV}(\{u\}) = \{u\}$ for any $u \in X$, $c^{IV}(\{x, y\}) = \{x, y\}$, $c^{IV}(\{y, z\}) = \{y, z\}$, $c^{IV}(\{x, z\}) = \{x, z\}$ and

$c^{IV}(\{x, y, z\}) = \{x, y\}$. Clearly, c^{IV} satisfies PR, C and SS. However, c^{IV} fails to satisfy CO since $z \in (c^{IV}(\{x, z\}) \cap c^{IV}(\{y, z\})) \setminus c^{IV}(\{x, y, z\})$;

(4) Let $X = \{x, y, z\}$, and $c^V \in C_X$ such that $c^V(\{u\}) = \{u\}$ for any $u \in X$, $c^V(\{x, y\}) = \{x\}$, $c^V(\{y, z\}) = \{y\}$, $c^V(\{x, z\}) = \{x, z\}$ and $c^V(\{x, y, z\}) = \{x\}$. Clearly, c^V satisfies PR, C and CO but fails to satisfy SS since $\emptyset \neq c^V(\{x, y, z\}) \subset c^V(\{x, z\})$.

Remark 16 Notice again that Theorem 14 above is essentially a refinement of well-known results due to Suzumura (see e.g. Suzumura (1983), Theorems 2.8 and 2.10) and Plott (1974), whose Theorems 3, 4, and 7 amount essentially to the equivalence between statements (iii), (iv) and (vii). It should also be mentioned here that the conjunction of C and SS turns out to be equivalent (see e.g. Suzumura (1983)) to another well-known and widely used property, namely

Path Independence (PI): for any $A, B \subseteq X$, $c(A \cup B) = c(c(A) \cup c(B))$

Thus, the equivalent statements of Theorem 14 are also equivalent to the statement ‘ $c \in C_X$ satisfies PR, PI and CO’.

It should be remarked that the characterizations provided above are in general quite straightforward extensions to *arbitrary* choice functions (with full domain) of previously known results concerning *proper* choice functions (with full domain). Indeed, the gist of the results offered in the present section may be summarized as follows:

(i) remarkably, the characterizations of *general* revealed cores and a-cores considered here consist of the very same properties used to characterize their nonempty-valued counterparts as supplemented with very mild-looking local nonemptiness requirements for choice sets of singleton and two-valued subsets, respectively;

(ii) the exact correspondence between revealed core-solutions and maximizing ‘rational’ choice functions is confirmed to hold within the general space of *arbitrary* choice functions: the alleged extra-generality of the latter subclass that has sometimes been alluded to in the literature (as e.g. in Suzumura (1983), p.21) does not materialize within the space of (total) choice functions and is therefore strictly confined to the realm of *partial* choice functions;

(iii) finally, and most notably, the class of general revealed cores turns out to inherit some of the supplementary order-theoretic structure enjoyed

by its larger ambient space as compared to the smaller and less regular space of proper choice functions: that is precisely the topic of the next section.

3 Posets and semilattices of revealed cores

Let us now turn to a global description of the order-theoretic structure of the class of all revealed core-solutions (a-core-solutions, nonempty-valued core-solutions, externally stable core-solutions, respectively).

A partially ordered set or *poset* is a pair $\mathbf{P} = (P, \leq)$ where P is a set and \leq is a reflexive, transitive and antisymmetric binary relation on P (i.e. for any $x \in P$, $x \leq x$ and for any $x, y, z \in P$, $x \leq z$ whenever $x \leq y$ and $y \leq z$, and $x = y$ whenever $x \leq y$ and $y \leq x$). For any $x \in P$ we posit $(x] = \{y \in P : y \leq x\}$. A *coatom* of a poset $\mathbf{P} = (P, \leq)$ with a top element or maximum 1_P is any $j \in P$ which is covered by 1_P - written $j < 1_P$ - i.e. $j < 1_P$ and $l = j$ for any $l \in P$ such that $j \leq l < 1_P$. The set of all coatoms of \mathbf{P} is denoted A_P^* . Dually, an *atom* of \mathbf{P} is any $j \in P$ which is an upper cover of 0_P - written $0_P < j$ - i.e. $0_P < j$ and $l = j$ for any $l \in P$ such that $0_P < l \leq j$. The set of all atoms of \mathbf{P} is denoted A_P .

A poset $\mathbf{P} = (P, \leq)$ is a *meet semilattice* (*join semilattice*, respectively) if for any $x, y \in P$ the \leq -greatest lower bound $x \wedge y$ (the \leq -least upper bound $x \vee y$, respectively) of $\{x, y\}$ does exist. Moreover, \mathbf{P} is a *lattice* if it is both a meet semilattice and a join semilattice.

A lattice $\mathbf{P} = (P, \leq)$ is *bounded* if there exist both a bottom element 0_P and a top element 1_P (hence in particular a finite lattice is also bounded), *distributive* iff $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$ for any $x, y, z \in P$, *complemented* if it is bounded and for any $x \in P$ there exists $x' \in P$ such that $x \vee x' = 1_P$ and $x \wedge x' = 0_P$, and *Boolean* iff it is both distributive and complemented.

A *meet semilattice* $\mathbf{P} = (P, \leq)$ is *lower distributive* if $((x], \leq_{(x)})$ is a distributive lattice for any $x \in P$, and has the *coronation* (or *join-Kelly*) *property* if -for any $x, y, z \in P$ - $((x \vee y) \vee z)$ exists in P whenever $x \vee y, x \vee z$ and $y \vee z$ also exist. A *meet semilattice* is *median* if it is *lower distributive* and has the *coronation property*.

The set C_X of all choice functions on X can be endowed in a natural way with the point-wise set inclusion partial order \leq by positing, for any $c, c' \in C_X$, $c \leq c'$ iff $c(A) \subseteq c'(A)$ for each $A \subseteq X$. Clearly, the identity operator c^{id} is its top element, and the constant empty-valued choice function c^\emptyset its bottom element. It is well-known, and easily checked, that (C_X, \leq) is

in fact a *Boolean lattice* with join $\vee = \cup$ (i.e. set-union) and meet $\wedge = \cap$ (i.e. set-intersection), both defined in the obvious component-wise manner: see e.g. Monjardet, Raderanirina (2004).

For any $x, y \in X$ such that $x \neq y$, $c_{xy}^+ \in C_X$ and $c_{xy}^- \in C_X$ are defined as follows: for all $A \subseteq X$, $c_{xy}^+(A) = A \setminus \{y\}$ if $\{x, y\} \subseteq A$, and $c_{xy}^+(A) = A$ otherwise, and $c_{xy}^-(\{z\}) = \{z\}$ for all $z \in X$, $c_{xy}^-(\{x, y\}) = \{y\}$, and $c_{xy}^-(A) = \emptyset$ for all $A \subseteq X$ such that $A \neq \{x, y\}$ and $\#A \neq 1$. Moreover, $C_+ = \{c_{xy}^+ : x, y \in X, x \neq y\}$, and $C_- = \{c_{xy}^- : x, y \in X, x \neq y\}$.

The *minimum ND choice function* $c^{[1]}$ is defined by the following rule: for any $x \in X$, $c^{[1]}(\{x\}) = \{x\}$, and $c^{[1]}(Y) = \emptyset$ for any $Y \subseteq X$ such that $\#Y \neq 1$.

Now, let $C_X^* \subseteq C_X$ denote the set of all revealed core-solutions on X , $C_X^{*a} \subseteq C_X^{*\circ}$ the set of all revealed asymmetric core-solutions, $C_X^{*\circ} = C_X^* \cap C_X^\circ$ the set of all revealed nonempty-valued core-solutions, and C_X^{*es} the set of all revealed externally stable core-solutions on X , respectively). We also denote with a slight abuse of notation (C_X^*, \leq) , (C_X^{*a}, \leq) , $(C_X^{*\circ}, \leq)$ and (C_X^{*es}, \leq) the corresponding subposets of (C_X, \leq) (where \leq denotes $\leq \cap (C_X^* \times C_X^*)$, $\leq \cap (C_X^{*a} \times C_X^{*a})$, $\leq \cap (C_X^{*\circ} \times C_X^{*\circ})$ and $\leq \cap (C_X^{*es} \times C_X^{*es})$, respectively). We have the following

Theorem 17 *The poset (C_X^*, \leq) of revealed core-solutions is a sub-meet-semilattice of (C_X, \leq) with c^{id} itself as its top element, but not a sub-join-semilattice of (C_X, \leq) . It also satisfies the coronation property hence it is a median meet semilattice. The bottom element of (C_X^*, \leq) is the minimum ND choice function $c^{[1]}$. Moreover, the set of coatoms of (C_X^*, \leq) is C_+ , and the set of its atoms is C_- .*

Proof. Let $c, c' \in C_X^*$, and consider $c \cap c'$. Clearly, for any $x \in X$, $(c \cap c')(\{x\}) = c(\{x\}) \cap c'(\{x\}) = \{x\}$ since c and c' satisfy ND: hence $c \cap c'$ does also satisfy ND.

Moreover, for any $A \subseteq B \subseteq X$, since c and c' both satisfy C, $(c \cap c')(B) \cap A = (c(B) \cap c'(B)) \cap A = c(B) \cap (c'(B) \cap A) \subseteq c(B) \cap c'(A) \subseteq c(B) \cap A \subseteq A$ hence $c \cap c'$ satisfies C.

Finally, since c and c' satisfy CO, for any $A, B \subseteq X$,

$$(c \cap c')(A) \cap (c \cap c')(B) =$$

$$(c(A) \cap c(B)) \cap (c'(A) \cap c'(B)) \subseteq c(A \cup B) \cap c'(A \cup B) = (c \cap c')(A \cup B)$$

and CO also holds for $c \cap c'$. It follows that, by Theorem 7 above, $c \cap c' \in C_X^*$,

whence (C_X^*, \leq) is a *sub-meet-semilattice* of (C_X, \leq) : in particular, it follows that (C_X^*, \leq) is *lower distributive*.

Furthermore, let us suppose that $c_1, c_2, c_3, c_1 \cup c_2, c_1 \cup c_3, c_2 \cup c_3 \in C_X^*$. Then, take $(c_1 \cup c_2) \cup c_3$ as defined in the obvious way. It is immediately checked that $(c_1 \cup c_2) \cup c_3$ does satisfy ND and C, by construction.

Thus, we only have to check that $(c_1 \cup c_2) \cup c_3$ does also satisfy CO. In order to check this last point, consider any $A, B \subseteq X$, and $x \in ((c_1 \cup c_2) \cup c_3)(A) \cap ((c_1 \cup c_2) \cup c_3)(B)$.

By definition, it follows that $x \in c_i(A) \cap c_j(B)$ for some $i, j = 1, 2, 3$. Hence, in particular, it also follows that $x \in (c_i \cup c_j)(A) \cap (c_i \cup c_j)(B)$ for some $i, j = 1, 2, 3$. Now, by hypothesis, $(c_i \cup c_j) \in C_X^*$ hence it satisfies CO. Therefore, $x \in (c_i \cup c_j)(A \cup B) \subseteq ((c_1 \cup c_2) \cup c_3)(A \cup B)$ and $(c_1 \cup c_2) \cup c_3$ also satisfies CO. As a consequence, $(c_1 \cup c_2) \cup c_3 \in C_X^*$: thus, (C_X^*, \leq) has the join-Kelly property and is therefore a median meet-semilattice as claimed.

It is easily checked that c^{id} , the top element of (C_X, \leq) , does also satisfy ND, C and CO hence as observed above $c \in C_X^*$ (see Example 2).

Now, consider $c^{[1]}$ as defined above: it satisfies ND, by definition, and, being nonempty-valued precisely on singletons, it trivially satisfies C and CO as well. Thus, $c^{[1]} \in C_X^*$. On the other hand, for any $c \in C_X^*$, c must satisfy ND, hence $c^{[1]} \leq c$.

Next, take any $c_{xy}^+ \in C_+$. Notice that, by definition, c_{xy}^+ satisfies ND. Also, if $A \subseteq B \subseteq X$ then the following cases may be distinguished: (a) $\{x, y\} \subseteq A$; (b) $\{x, y\} \not\subseteq A$ and $\{x, y\} \subseteq B$; (c) $\{x, y\} \not\subseteq B$. If $\{x, y\} \subseteq A$ then $c_{xy}^+(B) \cap A = A \setminus \{y\} = c_{xy}^+(A)$; if $\{x, y\} \not\subseteq A$ and $\{x, y\} \subseteq B$ then $c_{xy}^+(B) \cap A = (B \setminus \{y\}) \cap A = A \setminus \{y\} \subset A = c_{xy}^+(A)$; if $\{x, y\} \not\subseteq B$ then $c_{xy}^+(B) \cap A = A = c_{xy}^+(A)$: thus in any case C holds. Furthermore, let $z \notin c_{xy}^+(A \cup B)$: then by definition $z = y$ and $\{x, y\} \subseteq A \cup B$. Assume now that $y \in c_{xy}^+(A) \cap c_{xy}^+(B)$. Then, $\{x, y\} \not\subseteq A$ and $\{x, y\} \not\subseteq B$ while $y \in A \cap B$. It follows that $x \notin A \cup B$, a contradiction. Thus, CO is also satisfied by c_{xy}^+ , Theorem 7 applies, and $c_{xy}^+ \in C_X^*$.

Moreover, by definition $c_{xy}^+ < c^{id}$ i.e. $c_{xy}^+ \leq c^{id}$ and $c_{xy}^+ \neq c^{id}$.

Let $c \in C_X^*$ be such that $c_{xy}^+ \leq c \leq c^{id}$, and assume that $c_{xy}^+ \neq c$ i.e. there exists $A' \subseteq X$ such that $c_{xy}^+(A') \subset c(A') \subseteq A'$. Clearly, by Theorem 7, c satisfies ND, C and CO. If $c = c^{id}$ there is nothing to prove, so assume that there also exists $B \subseteq X$ such that $c_{xy}^+(B) \subseteq c(B) \subset B$. Notice that by definition of c_{xy}^+ , $c_{xy}^+(A') \subset A'$ entails $A' \supseteq \{x, y\}$ and $c_{xy}^+(A') = A' \setminus \{y\}$, hence in particular $c(A') = A'$. Also, there exists $z \in (B \setminus c(B)) \cap (B \setminus c_{xy}^+(B))$. By definition of c_{xy}^+ again, $z \in (B \setminus c_{xy}^+(B))$ entails $B \supseteq \{x, y\}$, $z = y$

and $c_{xy}^+(B) = B \setminus \{y\}$ (whence $x \in c_{xy}^+(B) \cap c(B)$). Therefore, $x \in c(B) \cap \{x, y\}$ whence, by C, $x \in c(\{x, y\})$.

Suppose first that $c(\{x, y\}) = \{x, y\}$, and consider $B \setminus \{x\}$. Clearly, by definition, $c_{xy}^+(B \setminus \{x\}) = c(B \setminus \{x\}) = B \setminus \{x\}$. Thus, $y \in c(\{x, y\}) \cap c(B \setminus \{x\})$ hence, by CO, $y \in c(B)$: a contradiction, since $y = z \in B \setminus c(B)$. Suppose then $c(\{x, y\}) = \{x\}$: since by hypothesis $c \in C_X^*$, there exists an irreflexive digraph (X, Δ) such that $c(A) = \mathbb{C}(A, \Delta_A)$ for any $A \subseteq X$. Therefore, $c(\{x, y\}) = \{x\}$ entails $x\Delta y$ that in turn entails $y \notin c(A')$ since $A' \supseteq \{x, y\}$: a contradiction again because $c(A') = A'$.

It follows that if $c_{xy}^+ \leq c \leq c^{id}$ then either $c = c^{id}$ or $c = c_{xy}^+$ i.e. c_{xy}^+ is indeed a *coatom* of (C_X^*, \leq) .

Conversely, let c be a coatom of (C_X^*, \leq) and suppose $c \notin C_+$. Then, for any pair of distinct $x, y \in X$, neither $c_{xy}^+ \leq c$ nor $c \leq c_{xy}^+$ i.e. there exist $A, B \subseteq X$ such that $c(A) \subset c_{xy}^+(A)$ and $c_{xy}^+(B) \subset c(B)$. Thus, by definition, $c_{xy}^+(B) = B \setminus \{y\}$ and $c(B) = B \supseteq \{x, y\}$, while there exists $z \in A$ such that $z \in c_{xy}^+(A) \setminus c(A)$. Hence, consider any $x \in A \setminus \{z\}$: then, there exists B' such that $\{x, z\} \subseteq B' \subseteq X$ and $c(B') = B'$. By C, $\{x, z\} = c(B') \cap \{x, z\} \subseteq c(\{x, z\})$ i.e. $c(\{x, z\}) = \{x, z\} \subseteq A$ for any $x \in A$ while $z \notin c(A)$, which contradicts CO in view of *finiteness* of X .

To check that each $c_{xy}^- \in C_-$ is an *atom* of (C_X^*, \leq) , notice first that $c_{xy}^- \in C_X^*$. Indeed, c_{xy}^- satisfies ND by construction. Also, if $A \subseteq B$ then $c_{xy}^-(B) \cap A \neq \emptyset$ entails that either $A = B = \{z\}$ for some $z \in X$, or $A \subseteq B \subseteq \{x, y\}$ i.e. either A is a singleton or $A = B$. Thus, in any case, if $A \subseteq B$ then by definition $c_{xy}^-(B) \cap A \subseteq c_{xy}^-(A)$ hence c_{xy}^- satisfies C. Moreover, for any $A, B \subseteq X$, if $x \in c_{xy}^-(A) \cap c_{xy}^-(B)$ then by definition of c_{xy}^- either $A = B = \{x\}$ or $(A \cup B) \in \{A, B\}$ and $A \cup B = \{x, y\}$: thus, in any case, $x \in c_{xy}^-(A \cup B)$ and CO is also satisfied by c_{xy}^- . Next, observe that $c_{xy}^-(A) = c^{[1]}(A)$ for any $A \neq \{x, y\}$, and $c_{xy}^-(\{x, y\}) = \{x\}$ while $c^{[1]}(\{x, y\}) = \emptyset$. Thus, for any $c \in C_X^*$ (indeed, for any $c \in C_X$) if $c^{[1]} \leq c \leq c_{xy}^-$ then either $c = c^{[1]}$ or $c = c_{xy}^-$.

Conversely, assume that c is an atom of (C_X^*, \leq) and $c \notin C_-$. Then, by definition of C_- , $c(A) = \emptyset$ for any A such that $\#A = 2$, and there exists $B \subseteq X$ such that $\#B \geq 3$ and $c(B) \neq \emptyset$. It follows that, for any $x \in c(B)$ and any $y \in B \setminus \{x\}$, $c(B) \cap \{x, y\} \not\subseteq \emptyset = c(\{x, y\})$, therefore violating C, a contradiction by Theorem 7.

To check that (C_X^*, \leq) is *not* a sub-join-semilattice of (C_X, \leq) , just consider without loss of generality $X = \{x, y, z\}$,

$R = \{(x, x), (y, y), (z, z), (y, x), (y, z), (x, z)\}$ and

$R^{-1} = \{(x, x), (y, y), (z, z), (x, y), (z, y), (z, x)\}$.

Now, posit $c_I(A) = \mathbb{C}(A, \Delta_A^R)$ and $c_{II}(A) = \mathbb{C}(X, \Delta_A^{R^{-1}})$ for any $A \subseteq X$. By definition $\mathbb{C}(X, \Delta^R) = \{y\}$, $\mathbb{C}(\{x, z\}, \Delta_{\{x, z\}}^R) = \{x\}$, $\mathbb{C}(X, \Delta^{R^{-1}}) = \{z\}$, and $\mathbb{C}(\{x, y\}, \Delta_{\{x, y\}}^{R^{-1}}) = \{x\}$ hence $(c_I \cup c_{II})(X) = \{y, z\}$, while $x \in (c_I \cup c_{II})(\{x, y\}) \cap (c_I \cup c_{II})(\{x, z\})$, which contradicts CO. ■

Remark 18 Notice that finiteness of X has been used in the proof above in order to show that the set of coatoms of (C_X^*, \leq) is contained in C^+ . The latter statement clearly holds for an infinite X as well provided CO is replaced with the following stronger version of ‘Concordance’

CO*: for any family $\{A_i\}_{i \in I}$ of subsets of X , $\bigcap_{i \in I} c(A_i) \subseteq c(\bigcup_{i \in I} A_i)$.

Remark 19 Since (C_X^*, \leq) is a semilattice with a top element (and indeed a finite one, under finiteness of X), it follows that it is also a lattice with meet $= \cap$ and join of a pair given by the meet of the (nonempty) set of upper bounds of that pair (see e.g. Davey, Priestley (1990)), which is however not a sublattice of (C_X, \leq) .

Thus, the poset of revealed core-solutions enjoys the remarkably regular structure of a *median meet-semilattice*. Notice that an important consequence of that fact is the following: any profile of revealed cores admits medians and the latter coincide with the *simple majority consensus* revealed core if the profile consists of an *odd* list of revealed cores. Therefore, in case *several revealed cores are to be considered for aggregation*, due perhaps to locally missing or unreliable data and/or plurality of information sources, an *amalgamation process by means of the simple majority aggregation rule is available* (see e.g. Monjardet, Raderanirina (2004) for some results on choice aggregation rules in the same vein).

The posets of revealed a-core-solutions, nonempty-valued core-solutions, and externally stable core-solutions are considerably less regular, as recorded by the following results, namely

Theorem 20 The poset (C_X^{*a}, \leq) of revealed a-core-solutions has a top element, c^{id} , and C_+ is the set of its coatoms, but it is neither a sub-meet-semilattice nor a sub-join-semilattice of (C_X, \leq) . The minimal elements of (C_X^{*a}, \leq) are the choice functions $c \in C_X$ that satisfy ND, 2-PR, C, CO and such that (a) $\#c(A) \leq 1$ for any $A \subseteq X$ and (b) not $D_{c'} \subset D_c$ for any c' that satisfies ND, 2-PR, C and CO.

Proof. To check that c^{id} is indeed the top element of (C_X^{*a}, \leq) it is only to be observed -in view of Theorem 7- that c^{id} does in fact also satisfy 2-PR. Similarly -in view of Theorem 7 and of the proof of Theorem 17 provided above- to see that C_+ is the set of coatoms of (C_X^{*a}, \leq) it is only to be checked that any $c_{xy}^+ \in C_+$ does also satisfy 2-PR (which is clearly the case, by definition).

The proof of Theorem 17 already establishes that (C_X^{*a}, \leq) is not a sub-join-semilattice of (C_X, \leq) since, as it is easily checked, c_I and c_{II} as defined there do belong to C_X^{*a} .

Next, consider c_{III} and c_{IV} defined as follows: assume without loss of generality $X = \{x, y, z\}$, and take $\Delta^{III} = \{(x, y), (x, z), (y, z)\}$, $\Delta^{IV} = \{(x, y), (x, z), (z, y)\}$ (notice that both (X, Δ^{III}) and (X, Δ^{IV}) are asymmetric digraphs); then, for any $A \subseteq X$, posit $c_{III}(A) = \mathbb{C}(A, \Delta_A^{III})$ and $c_{IV}(A) = \mathbb{C}(A, \Delta_A^{IV})$. Clearly, by definition, $\{c^{III}, c^{IV}\} \subseteq C_X^{*a}$.

However, $(c^{III} \cap c^{IV})(\{y, z\}) = \mathbb{C}(\{y, z\}, \Delta_{\{y, z\}}^{III}) \cap \mathbb{C}(\{y, z\}, \Delta_{\{y, z\}}^{IV}) = \{y\} \cap \{z\} = \emptyset$.

Therefore, $c^{III} \cap c^{IV}$ violates 2-PR hence by Theorem 7 $c^{III} \cap c^{IV} \notin C_X^{*a}$. It follows that (C_X^{*a}, \leq) is not a sub-meet-semilattice of (C_X, \leq) .

The last statement about minimal elements of (C_X^{*a}, \leq) is a straightforward consequence of Theorem 10. ■

Theorem 21 *The poset $(C_X^{*\circ}, \leq)$ of nonempty-valued core-solutions has a top element, c^{id} , and C_+ is the set of its coatoms, but it is neither a sub-meet-semilattice nor a sub-join-semilattice of (C_X, \leq) . The minimal elements of (C_X^{*a}, \leq) are the single-valued choice functions that satisfy C and CO.*

Proof. First, notice that by definition c^{id} is proper, hence $c^{id} \in C_X^{*\circ}$ since as previously shown it is a core-solution. Also, it is immediately checked that, by definition, any c_{xy}^+ is *proper*. Therefore, the proof of Theorem 17 also establishes that C_+ is the set of coatoms of $(C_X^{*\circ}, \leq)$. In the same vein, it is immediately checked that $c_I, c_{II}, c_{III}, c_{IV}$ -as defined above in the proofs of the two previous Theorems- are also proper. It follows, by those proofs, that $(C_X^{*\circ}, \leq)$ is neither a sub-meet-semilattice nor a sub-join-semilattice of (C_X, \leq) . The final statement about minimal elements of $(C_X^{*\circ}, \leq)$ is an immediate consequence of Corollary 12. ■

Theorem 22 *The poset (C_X^{*es}, \leq) of revealed externally stable core-solutions, has a top element, c^{id} , and C_+ is the set of its coatoms, but it is neither a sub-*

meet-semilattice nor a sub-join-semilattice of (C_X, \leq) . The minimal elements of (C_X^{*a}, \leq) are the single-valued choice functions that satisfy C, CO and SS.

Proof. Observe that for any $A \subseteq B \subseteq X$, if $c^{id}(B) \subseteq c^{id}(A)$ then of course $B \subseteq A$ i.e. $B = A$ whence $c^{id}(A) = c^{id}(B)$ and SS is clearly satisfied by c^{id} . In view of Theorem 14, this establishes that c^{id} is also the top element of (C_X^{*es}, \leq) . Also, it is immediately checked that any c_{xy}^+ satisfies SS: indeed, let $A, B \subseteq X$ be such that $A \subseteq B$ and $\emptyset \neq c_{xy}^+(B) \subseteq c_{xy}^+(A)$. Since $A \subseteq B$, the following jointly exhaustive cases are to be distinguished: (a) $\{x, y\} \subseteq A \cap B$; (b) $\{x, y\} \not\subseteq A \cup B$; (c) $\{x, y\} \subseteq B$ and $\{x, y\} \not\subseteq A$. Under (a), $c_{xy}^+(A) = A \setminus \{y\}$ and $c_{xy}^+(B) = B \setminus \{y\}$ hence $c_{xy}^+(A) \subseteq c_{xy}^+(B)$. Under (b), $c_{xy}^+(A) = A$ and $c_{xy}^+(B) = B$ hence again $c_{xy}^+(A) \subseteq c_{xy}^+(B)$. Under (c), $c_{xy}^+(A) = A$ and $c_{xy}^+(B) = B \setminus \{y\}$ whence $A \neq B$ i.e. $A \subset B$. By hypothesis, $c_{xy}^+(B) \subseteq c_{xy}^+(A)$ hence $B \setminus \{y\} \subseteq A \subset B$: thus, $y \notin A$ and $B = A \cup \{y\}$ and therefore $c_{xy}^+(B) = B \setminus \{y\} = A = c_{xy}^+(A)$. It follows that c_{xy}^+ does in fact satisfy SS. Therefore, the proof of Theorem 17 also establishes that C_+ is the set of coatoms of (C_X^{*es}, \leq) .

Finally, it is immediately checked by direct inspection that $c_I, c_{II}, c_{III}, c_{IV}$ -as defined above in the proofs of Theorems 19 and 20- do also (trivially) satisfy SS. It follows, by the very same proofs, that (C_X^{*es}, \leq) is neither a submeet-semilattice nor a sub-join-semilattice of (C_X, \leq) . The final statement about minimal elements of (C_X^{*es}, \leq) is an immediate consequence of Theorem 14. ■

Thus, while only the poset of revealed core-solutions is a (meet) sub-semilattice of (C_X, \leq) all the posets of revealed cores defined above share their top element and set of coatoms.

4 Concluding remarks

Choice functions with full domain which may be regarded as core-solutions or externally stable core solutions of an underlying dominance digraph (X, Δ) have been characterized both in the general case and for asymmetric dominance digraphs. Both characterizations combine a version of the usual mix of contraction consistency and expansion consistency conditions which is required for the special case of proper i.e. nonempty-valued choice functions with a suitable local nonemptiness requirement for choice sets. The characterizations provided above have also been shown to be helpful for a simple

analysis of the basic order-theoretic structure of revealed cores. In particular, as mentioned in the Introduction, every revealed core embodies a considerable part of the structure of standard maximizing choice functions, while the global structure of (full domain) revealed cores retains precisely the median semi-latticial properties of the space of all (full domain) choice functions that are most significant from the point of view of simple majority aggregation. The latter property, however, is not shared by asymmetric or externally stable revealed cores.

References

- [1] Aizerman M., Aleskerov F. (1995): *Theory of choice*. North Holland: Amsterdam.
- [2] Bossert W., Suzumura K. (2010): *Consistency, choice, and rationality*. Harvard University Press: Cambridge MASS.
- [3] Danilov V., Koshevoy G. (2009): Choice functions and extensive operators, *Order* 26, 69-94.
- [4] Davey B.A., Priestley H.A. (1990): *Introduction to lattices and order*. Cambridge University Press: Cambridge UK.
- [5] Monjardet B. (2007): Some order dualities in logic, games and choices, *International Game Theory Review* 9, 1-12.
- [6] Monjardet B., Raderanirina V. (2004): Lattices of choice functions and consensus problems, *Social Choice and Welfare* 23, 349-382.
- [7] Plott C. (1974): On game solutions and revealed preference theory, *Social Science Working Paper 35*, California Institute of Technology, Division of the Humanities and Social Sciences, Pasadena CA.
- [8] Sen A.K. (1971): Choice functions and revealed preference, *Review of Economic Studies* 38, 307-317.
- [9] Suzumura K. (1983): *Rational choice, collective decisions, and social welfare*. Cambridge University Press: Cambridge UK.

- [10] Vannucci S. (2009): Choosing VNM-stable sets of the revealed dominance digraph, DEP Working Paper 576, Siena.
- [11] Wilson R.B. (1970): The finer structure of revealed preference, *Journal of Economic Theory* 2, 348-353.